



## **GUIDELINE ONLY - Equine Vaccine Schedule**

Vaccine	Protects Against	Time to give	Booster needed?	
Encevac-T with Havolgen (an adjuvant used to create a better immune response) or Equiloid	EEE, WEE, and tetanus	Beginning of mosquito season (April)	Needs a booster 3-4 weeks later if first dose. If vaccinating once yearly, should be given late spring (end of May or early June). If given early spring, booster should be given in the fall.	
Fluvac Innovator 5	EEE, WEE, tetanus, rhino, & influenza	Beginning of mosquito season (April)	Should booster in the fall for a full year's protection. Needs a booster 3-4 weeks if first dose. If vaccinating once yearly, should be given late spring (end of May or early June).	
Vetera 5XP	EEE, WEE, tetanus, rhino, & influenza	Beginning of mosquito season (April)	Should booster in the fall for a full year's protection. Needs a booster 3-4 weeks if first dose. If vaccinating once yearly, should be given late spring (end of May or early June).	
West Nile Innovator	West Nile	Beginning of mosquito season	Needs booster in 3-4 weeks if first dose, then annually.	
Tetanus Toxoid	Tetanus	Anytime	Needs booster in 3-4 weeks if first dose. If cut, booster required if vaccine has not been given within 6 months.	



## GUIDELINE ONLY - Equine Vaccine Schedule

Pinnacle	Strangles	Anytime; give vaccine by itself, separate by 2 days	Needs booster in 2-3 weeks if first dose, then every 6 months.	**Given intranasal – very important
West Nile Innovator and EWT	West Nile, WEE & EEE	Beginning of mosquito season (April)	Should do fall EWT booster. Needs booster in 3-4 weeks if first dose. If vaccinating once yearly, should be given late spring (end of May or early June).	
Pneumabort-K +1b	Rhinopneumonitis			Given to breeding mares
Botulism (Botvax B)	Botulism	Annually	If first dose, then 2nd booster is needed 4 weeks after 1st dose. 3rd booster required weeks after 2nd dose.	
Imrab	Rabies (not typically done in this area, but available)	Annually	No, single does annually	May be given to mares prior to breeding to reduce number given during pregnancy
Equine Potomavac & Imrab	Potomac Horse Fever (PHF)	Annually	Booster needed 3-4 weeks after initial if first dose	May re-vaccinate 3-4 months in areas where risk is high